

See discussions, stats, and author profiles for this publication at: <http://www.researchgate.net/publication/263934329>

The Potential Therapeutic Effects of THC on Alzheimer's Disease

ARTICLE *in* JOURNAL OF ALZHEIMER'S DISEASE: JAD · JULY 2014

Impact Factor: 4.15 · DOI: 10.3233/JAD-140093 · Source: PubMed

CITATIONS

5

READS

1,203

9 AUTHORS, INCLUDING:



[Yaqiong Li](#)

University of South Florida

15 PUBLICATIONS 79 CITATIONS

[SEE PROFILE](#)



[Xiaoyang Lin](#)

University of South Florida

35 PUBLICATIONS 724 CITATIONS

[SEE PROFILE](#)



[Neel Nabar](#)

National Institute of Allergy and Infectious...

8 PUBLICATIONS 28 CITATIONS

[SEE PROFILE](#)



[Jianfeng Cai](#)

University of South Florida

70 PUBLICATIONS 835 CITATIONS

[SEE PROFILE](#)

The Potential Therapeutic Effects of THC on Alzheimer's Disease

Chuanhai Cao^{a,b,*}, Yaqiong Li^c, Hui Liu^{a,b}, Ge Bai^c, Jonathan Mayl^b, Xiaoyang Lin^{a,b}, Kyle Sutherland^d, Neel Nabar^e and Jianfeng Cai^{c,*}

^aCollege of Pharmacy, University of South Florida, Tampa FL, USA

^bUSF-Health Byrd Alzheimer's Institute, University of South Florida, Tampa FL, USA

^cDepartment of Chemistry, University of South Florida, Tampa FL, USA

^dCollege of Medicine, University of South Florida, Tampa FL, USA

^eThomas Jefferson University, Philadelphia, PA, USA

Accepted 29 April 2014

Abstract. The purpose of this study was to investigate the potential therapeutic qualities of Δ^9 -tetrahydrocannabinol (THC) with respect to slowing or halting the hallmark characteristics of Alzheimer's disease. N2a-variant amyloid- β protein precursor (A β PP) cells were incubated with THC and assayed for amyloid- β (A β) levels at the 6-, 24-, and 48-hour time marks. THC was also tested for synergy with caffeine, in respect to the reduction of the A β level in N2a/A β PPswe cells. THC was also tested to determine if multiple treatments were beneficial. The MTT assay was performed to test the toxicity of THC. Thioflavin T assays and western blots were performed to test the direct anti-A β aggregation significance of THC. Lastly, THC was tested to determine its effects on glycogen synthase kinase-3 β (GSK-3 β) and related signaling pathways. From the results, we have discovered THC to be effective at lowering A β levels in N2a/A β PPswe cells at extremely low concentrations in a dose-dependent manner. However, no additive effect was found by combining caffeine and THC together. We did discover that THC directly interacts with A β peptide, thereby inhibiting aggregation. Furthermore, THC was effective at lowering both total GSK-3 β levels and phosphorylated GSK-3 β in a dose-dependent manner at low concentrations. At the treatment concentrations, no toxicity was observed and the CB1 receptor was not significantly upregulated. Additionally, low doses of THC can enhance mitochondria function and does not inhibit melatonin's enhancement of mitochondria function. These sets of data strongly suggest that THC could be a potential therapeutic treatment option for Alzheimer's disease through multiple functions and pathways.

Keywords: Alzheimer's disease, amyloid- β peptide, cannabinoid, CB1 receptor, CB2 receptor, delta(9)-tetrahydrocannabinol, neurodegeneration

INTRODUCTION

In 2011 alone, 15 million family members have provided more than 17.4 billion hours of care to diagnosed Alzheimer's disease (AD) patients. That care translates into more than \$210 billion of AD-related services [1]. This disease translates into an enormous burden on caregivers, as well as the health care system, both med-

ically and economically. To date, there have been no effective treatments developed to cure or delay the progression of AD [2, 3]. By 2050, an estimated 11 to 16 million Americans will be living with the disease [1, 4].

AD pathology can be divided into two categories, familial inherited AD and sporadic AD. The histopathologies of early onset familial AD and late onset sporadic AD are indistinguishable. Both forms of AD are characterized by extracellular amyloid- β (A β) peptide, and by amyloid plaques and tau-containing neurofibrillary tangles [3]. The misfolded structure of the A β peptides generates a characteristic tendency for their aggregation [5]. It has long been believed

*Correspondence to: Chuanhai Cao, PhD, College of Pharmacy, University of South Florida, USF-Health Byrd Alzheimer's Institute, 4001 E. Fletcher Avenue, Tampa, FL 33613, USA. Tel.: +1 813 3960742; Email: ccao@health.usf.edu and Jianfeng Cai, PhD, Department of Chemistry, University of South Florida, Tampa, FL 33620, USA. E-mail: jianfengcai@usf.edu.

47 that $A\beta_{1-40}$ ($A\beta_{40}$) and $A\beta_{1-42}$ ($A\beta_{42}$) aggregates are
48 the constituents of the insoluble plaques that are char-
49 acteristic of AD. This disease is also associated with
50 neuroinflammation, excitotoxicity, and oxidative stress
51 [6, 7]. However, the continuous aggregation of $A\beta$
52 peptides along with hyperphosphorylation of the tau
53 protein inside the cell, causing neurofibrillary tangle
54 formation, are generally accepted as the major etiolog-
55 ical factors of the neuronal cell death associated with
56 the progression of AD [8–10].

57 Recent studies have also suggested that glycogen
58 synthase kinase 3 (GSK-3) has a key role in the patho-
59 genesis of both sporadic and familial AD [11, 12]. It
60 has been reported that GSK-3 β induces hyperphos-
61 phosphorylation of tau [13–17]. Moreover, overexpression
62 of GSK-3 in Tet/GSK-3 β mice reveal pathological
63 symptoms that correspond to AD pathology with
64 respect to spatial learning deficits, reactive astrocyto-
65 sis, increased $A\beta$ production, and plaque associated
66 inflammation, as well as tau hyperphosphorylation
67 resulting in $A\beta$ -mediated neuronal death [18]. Addi-
68 tionally, chronic lithium (GSK-3 inhibitor) treatment
69 in double transgenic mice overexpressing GSK-3 β and
70 tau has shown to prevent tau hyperphosphorylation
71 and neurofibrillary tangle formation [19]. Some reports
72 have also indicated that GSK-3 α plays a role in regu-
73 lating amyloid- β protein precursor ($A\beta$ PP) cleavage,
74 resulting in increased $A\beta$ production [20, 21]. It has
75 also been shown that the $A\beta$ load in mouse brain can
76 be robustly ameliorated by the inhibition of GSK-3 β
77 [22].

78 Along with past research suggesting an involve-
79 ment of GSK-3 in the pathogenesis of AD, there
80 have also been recent studies suggesting the intricate
81 involvement of the cannabinoid system in AD. It was
82 reported that the cannabinoid system can limit the neu-
83 rodegenerative processes that drive the progression of
84 the disease, and may provide a new avenue for dis-
85 ease control [23]. Currently the complete pathway and
86 mechanism of action of the cannabinoid system are
87 unknown, however, studies have been conducted to
88 determine the involvement of the cannabinoid 1 (CB_1)
89 and cannabinoid 2 (CB_2) receptors in AD brain [6]. The
90 CB_1 receptor is abundant in the brain and contributes to
91 learning, memory, and cognitive processes which are
92 interrupted early in the course of AD [24]. To the con-
93 trary, CB_2 receptor expression is more limited and has
94 been anatomically found in neurons within the brain-
95 stem [25], cerebellum [26], and microglia [27]. Recent
96 research has also investigated the propensity of endo-
97 cannabinoid receptor sub-types 1 (CB_1) and 2 (CB_2)
98 to elicit a neuroprotective and anti-inflammatory effect

99 on the brain when stimulated by endocannabinoids
100 [28]. Postmortem studies of AD brains have detected
101 increased expression of CB_1 and CB_2 receptors on
102 microglia within the plaque, while CB_1 expression is
103 reduced in neurons more remote from the plaque [29].
104 It is also noted that the endocannabinoid metabolizing
105 enzyme, fatty acid amide hydrolase, is upregulated in
106 the plaque [30]. There is also an increase in expres-
107 sion of anandamide metabolites, such as arachidonic
108 acid, in the vicinity of the plaque [30]. These find-
109 ings may indirectly suggest that the increase in CB_1
110 and CB_2 receptors may be to offset the lack of activity
111 with their ligands due to increased metabolic activ-
112 ity of fatty acid amide hydrolase. These alterations
113 in the cannabinoid system suggest an involvement of
114 endogenous cannabinoids in the pathogenesis of AD
115 or that this system may be altered by the pathophysiol-
116 ogy of the disease [6]. Understanding that microglial
117 activation is reserved in all cases of AD, it is important
118 to identify that endogenous cannabinoids prevent $A\beta$ -
119 induced microglial activation both *in vitro* and *in vivo*
120 [31]. These receptors are known to experience time
121 dependent and brain region specific alterations during
122 neurodegenerative and neuroinflammatory disorders to
123 attempt to counteract excitotoxicity and inflammation
124 [32].

125 Endocannabinoid receptors, CB_1 and CB_2 , have
126 been reported to interact with the endocannabinoid
127 molecules: 2-arachidonoyl glycerol and anandamide.
128 However, it has also been reported that CB_1 and
129 CB_2 also react interact with Δ^9 -tetrahydrocannabinol
130 (THC) isolated from the *Cannabis sativa* plant [33].
131 Furthermore, early reports indicate that Dronabinol,
132 an oil-based solution of Δ^9 -THC, improves the dis-
133 turbed behavior and stimulates appetite in AD patients
134 [34], and alleviates nocturnal agitation in severely
135 demented patients [35]. Accumulated evidence also
136 suggests antioxidants having anti-inflammatory and
137 neuroprotective roles [23].

138 It has also been shown that THC can decrease
139 the level of $A\beta$ -induced increases in reactive oxygen
140 species, decreases in mitochondrial membrane poten-
141 tial, and caspase (a protein that is intimately involved
142 in the regulation of apoptosis) activation, as well as
143 protect human neurons from oligomeric $A\beta$ -induced
144 toxicity [36]. While it is understood that cannabi-
145 noids are active against inflammation, our research
146 investigated the neuroprotective properties of THC,
147 the active component of marijuana. Here we evalu-
148 ated: 1) the effects of THC against $A\beta$ expression in
149 N2a/ $A\beta$ PPsw cells against the effects of caffeine, a
150 reported $A\beta$ expression suppressor [37]; 2) the direct

151 effects of THC against A β aggregation, one patho-
152 logical marker of AD; 3) the mechanism behind the
153 anti-pathological properties of THC on AD; 4) the tox-
154 icity of THC and caffeine individually; and 5) the effects
155 of THC on GSK-3 β and other related signal pathways
156 in N2a/A β PPswe cells.

157 MATERIALS AND METHODS

158 *Drugs used in this study*

159 THC solution was purchased from Sigma (T4764-
160 1ML Sigma Aldrich); caffeine was purchased from
161 Sigma (C0750-100G, Sigma Aldrich); melatonin was
162 purchased from Sigma (M5250-5G, Sigma Aldrich).

163 *ELISA for detection of total A β in protein samples*

164 50 μ l of goat anti-PWT1-42 antibody solution was
165 added to the sample and incubated overnight, followed
166 by a 1-hour incubation with 0.1% I-block buffer. The
167 tissue culture supernatant was diluted 1:10 with dilu-
168 ent buffer containing a protease inhibitor. Standards
169 (1000, 500, 250, 125, 62.5, 31.25 pg/ml) were prepared
170 by serial dilution. The plate was washed and 50 μ l of
171 sample or standard was added with triplication. 50 μ l
172 of both Biosource 40/42 (HS) (primary antibody) A β
173 and a standard solution was added to each well and
174 incubated for 3 hours followed by 5 \times wash with PBST.
175 100 μ l prepared secondary antibody (1:350 anti-rabbit
176 HRP) was added and incubated at 37 $^{\circ}$ C for 45 minutes
177 on a shaker. The plate was washed; TMB substrate
178 was added (100 μ l) and incubated for 10–30 minutes
179 in the dark. The reaction was halted by adding 100 μ l
180 stop solution for detection at 450 nm. A 4 parameter
181 regression was used for the standard.

182 *Cell culture and drug treatment*

183 N2a/A β PPswe cells, N2a cells stably expressing
184 human A β PP carrying the K670N/M671L Swedish
185 mutation (A β PPswe), were grown in Dulbecco's modi-
186 fied Eagle medium containing 10% fetal bovine serum,
187 100 U/mL penicillin, 100 μ g/mL streptomycin, and
188 400 μ g/mL G418 (Invitrogen), at 37 $^{\circ}$ C in the pres-
189 ence of 5% CO $_2$. N2a/A β PPswe cells were diluted
190 with medium to a concentration of 2 \times 10 5 /ml, and
191 plated into the each well in 3 ml. 2 ml of trypsin was
192 incubated at room temperature, or 37 $^{\circ}$ C. When most of
193 the cells began to float, trypsin was decanted and 5 ml
194 of fresh pre-warmed medium was added. Pipetting was
195 performed more than 30 times to ensure cells were sep-

196 arated into individual cells. One drop of medium was
197 put into 1.5 ml tubes for counting; 10 μ l of trypan blue
198 and 10 μ l of medium of cells were added and applied to
199 cytometer for counting. The rule was total number of
200 cells of all for diagonal blocks/4 X 2 X 10000 = number
201 of cells/ml. The proper amount of cell medium and
202 fresh medium was added into new flasks according
203 to the ratio of dilution. Pipetting was performed 10
204 times to homogenize cells. 3 ml of cells were seeded
205 into medium into each 6 well plate. When one pipette
206 was used up, the cells were mixed in the flask before
207 using them for the next pipette. Compounds for screen-
208 ing were resolved in DMSO, at 1000 fold to the final
209 concentration in the well. Pipetting of 10 μ l solution,
210 then addition into 990 μ l medium was performed; mix-
211 ing followed. 12 hours after cells were plated, 400 μ l
212 of compounds were added into 3.6 ml medium. The
213 medium was then removed from the six-wells. 3 ml of
214 medium with 1% DMSO was added to well 1; in well
215 2, 3 ml melatonin solution was added. In well 3, 4, 5,
216 and 6, compound solutions of 3 ml were added.

217 *MTT assay*

218 Cells were plated in 96-well tissue culture plate
219 at 10,000 cells/well, 100 μ l/well. 100 μ l THC solu-
220 tion was added at 2 \times concentrations in each well.
221 Control groups are: 1) cells without THC treatment,
222 cells and fresh medium only; and 2) blank, wells with
223 medium without cells. All wells were replicated. Wells
224 were incubated for 36 hours. Cell proliferation kit
225 (Roche 11465007001) was then applied for toxicity
226 assay according to the standard protocol. 10 μ l of MTT
227 reagent was first added to each well and incubated at
228 37 $^{\circ}$ C for 4 hours. Then 100 μ l of solubilization solu-
229 tion was added to each well. These were incubated
230 overnight and optical density (OD) values were read at
231 575 nm. The percentage of cell viability was calculated
232 as: Cell viability% = (OD – OD blank) / (OD control –
233 OD blank)

234 *Western blot for anti-aggregation assay*

235 HFIP pretreated A β $_{1-40}$ peptide were obtained
236 from Biomer Technology, California. A β $_{1-40}$ pep-
237 tide solution was prepared in Ham's F-12 solution to
238 concentrations of 200 μ M as stock. In the 15 μ l aggre-
239 gation system: 1) THC at final concentration of 25
240 nM, 2.5 nM, or 0.25 nM; and 2) 1.5 μ l peptide stock
241 solution was added. Then 15 μ l with F-12 medium
242 was made. Aggregation was allowed for 48 hours at
243 37 $^{\circ}$ C. After incubation, isomers of A β peptide were

244 separated by 12% Tris-Tricine gel electrophoresis at
 245 100 V for 180–210 minutes, a temperatures under 4°C.
 246 The protein was transferred to PVDF membrane with
 247 semi-dry transfer at 200 mA for 70 minutes. For west-
 248 ern blot detection, the membrane was blocked with
 249 1.5% BSA solution in PBST solution (0.5%), then
 250 incubated in 1st Antibody: 6E10 (Signet) 1 mg/ml,
 251 diluted by 1:1000 dilution in blocking buffer. It was
 252 then washed 3 times with 1× PBST solution, and
 253 then incubated with second antibody (anti-mouse IgG-
 254 HRP sigma A9044. 1:5000 diluted in blocking buffer).
 255 After membrane was developed, film with bands were
 256 scanned, followed by analysis of gel-quantification
 257 software (QuantityOne, from Bio-rad).

258 *Thioflavin T fluorescence assay*

259 HFIP pretreated A β_{1-40} peptide was obtained from
 260 Biomer Technology, California. In thioflavin T (ThT)
 261 solutions (1.6 μ g/ml dissolved in 20 mM Tris-HCL),
 262 THC solution was prepared at concentration of 250,
 263 25, 2.5, and 0.25 nM. THC solution was added contain-
 264 ing ThT buffer into black 96 well plates. Unaggregated
 265 A β peptide solution was thawed, diluted, and imme-
 266 diately added to wells, making the final concentration
 267 of A β_{1-40} at 1 μ M. Control groups were setup as: 1)
 268 aggregation control; 2) control with ThT buffer only;
 269 and 3) Tris-HCl buffer only. Plate was mixed and flu-
 270 orescence was read at 482 nm with excitation 440 nm
 271 with Biotek All-in-One plate reader. Fluorescence was
 272 screened for 2 hours with 5-minute intervals.

273 *Western blot for total and phosphorylated GSK-3 β , 274 total tau and phosphorylated tau, and β -actin*

275 Followed by THC treatment in tissue culture,
 276 N2a/A β PPswe cell lysate were collected, quantified,
 277 and aliquoted. Using 12% Tris-Glycine gel system
 278 (Biorad), protein were separated by electrophoresis
 279 and semi-dry transferred to PVDF membrane. GSK-3 β
 280 and β -actin antibodies were used as primary antibody.
 281 After adding secondary antibody, the membranes were
 282 exposed using ECL substrate (Pierce). After mem-
 283 brane was developed, film with bands were scanned,
 284 followed by analysis of gel-quantification software
 285 (QuantityOne, from Biorad).

286 *Mitochondria isolation and respiratory 287 measurements*

288 The respiratory function of isolated mitochondria
 289 was measured using a miniature Clark type oxygen

290 electrode (Strathkelvin Instruments, MT200A cham-
 291 ber, Glasgow, UK). Detail method is published in
 292 Dragicevic et al. [38].

293 *Statistical analysis and graphs*

294 All data were analyzed with one-way ANOVA and
 295 *post hoc* analysis was conducted with Turkey's group
 296 analysis and $p < 0.05$ was considered as statistical sig-
 297 nificance (GraphPad 6.0). All graphs were graphed
 298 with GraphPad 6.0 software.

299 **RESULTS**

300 *THC can decrease A β level in N2a/A β PPswe*

301 ELISA assay was performed for A β_{40} levels in
 302 N2a/A β PPswe cells 6 hours after cells were treated
 303 at different concentrations individually with THC, and
 304 caffeine—a reported compound to lower serum A β_{40}
 305 levels in a mouse model [39]—showed a significant
 306 reduction in A β_{40} levels of THC and caffeine ver-
 307 sus the control (Fig. 1A). However, 24 hours after
 308 treatment of N2a/A β PPswe cells, A β_{40} concentra-
 309 tions were measured again in the THC treated cells
 310 versus the control. An increasing difference in A β_{40}
 311 concentrations were noted in both THC treated cells
 312 and caffeine treated cells in a dose-dependent man-
 313 ner (Fig. 1B). The assay was performed again, 48
 314 hours after treatment of N2a/A β PPswe cells with THC
 315 versus the control at each concentration of the drugs
 316 originally used. THC-treated N2a/A β PPswe cells sig-
 317 nificantly differed more in A β_{40} concentrations versus
 318 the control then at the 6- and 24-hour time point. The
 319 significant difference was conserved and greater over
 320 each increasing dose of THC and caffeine adminis-
 321 tered versus the control (Fig. 1C). These data suggest
 322 THC's and caffeine's inherent anti-A β_{40} properties are
 323 time and dose dependent in N2a/A β PPswe cell mod-
 324 els. This data also reveals that THC may delay or halt
 325 the progression of AD by inhibiting the production of
 326 A β_{40} peptide in the central nervous system.

327 *Synergy between THC and caffeine on A β_{40} 328 concentration in N2a/A β PPswe cells*

329 THC and caffeine were assayed for a synergistic
 330 effect on A β_{40} concentration in N2a/A β PPswe cells
 331 (Fig. 2). However, no synergistic properties of THC
 332 and caffeine are seen as there is no significant differ-
 333 ence in the concentration of A β_{40} in N2a/A β PPswe
 334 cells solely treated with THC as compared to cells




Fig. 1. (A) $A\beta_{40}$ (pg/ml) *in vitro* measured 6 hours from incubation in N2a/A β PPsw cells. Three groups of cells were assayed: 1) those that were not treated with THC; 2) those that were treated with THC; and 3) those that were treated with caffeine. Treatment in both the THC group and in the caffeine group resulted in a dose-dependent decrease in $A\beta_{40}$ concentration after 6 hours. There are no significant differences among all groups ($p > 0.05$). The concentrations of THC from A to F are 0 nM, 0.25 nM, 2.5 nM, 25 nM, 250 nM, and 2500 nM respectively, and concentrations of caffeine from A to F are 0 μ M, 0.625 μ M, 1.25 μ M, 2.5 μ M, 5 μ M, and 10 μ M, respectively. (B) $A\beta_{40}$ (pg/ml) *in vitro* measured 24 hours from incubation in N2a/A β PPsw cells. A dose-dependent decrease in concentration of $A\beta_{40}$ was still observed. THC: A, B, C, versus F are $p < 0.05$ and all other groups in comparison are $p > 0.05$. Caffeine: A versus B, B versus E, and all other groups versus F are $p < 0.05$. The concentrations of THC from A to F are 0, 0.25 nM, 2.5 nM, 25 nM, 250 nM, and 2500 nM, respectively, and concentrations of caffeine from A to F are 0, 0.625 μ M, 1.25 μ M, 2.5 μ M, 5 μ M, and 10 μ M, respectively. (C) $A\beta_{40}$ (pg/ml) *in vitro* measured 48 hours from incubation in N2a/A β PPsw cells. A dose-dependent decrease in $A\beta_{40}$ (pg/ml) in conserved. THC groups: $p > 0.05$ for A versus B, and all other groups are $p < 0.05$. Caffeine groups: $p < 0.05$ for B versus D, and all other comparisons between groups are $p > 0.05$. The concentrations of THC from A to F are 0, 0.25 nM, 2.5 nM, 25 nM, 250 nM, and 2500 nM, respectively, and concentrations of caffeine from A to F are 0, 0.625 μ M, 1.25 μ M, 2.5 μ M, 5 μ M, and 10 μ M, respectively.

355 treated with 2.5 μ M caffeine and THC at various
356 concentrations.

357 *Repeated treatment can continuously decrease A β*
358 *production*

359 Our data also illustrates N2a/A β PPsw cells treated with
360 THC twice, 24 hours apart from each treatment,
361 showed a significant decrease in $A\beta_{40}$ concentration
362 compared to cells treated once (Fig. 3A). While
363 the decrease in $A\beta_{40}$ expression is not observed
364 at concentration close to 10 μ M, they are seen at
365 25 μ M and greater suggesting multiple treatments
366 may be efficacious in reducing $A\beta_{40}$ concentration in
367 N2a/A β PPsw cells and animal models.

368 *Cell toxicity detection of THC on N2a/A β PPsw*
369 *cells*

370 THC was also measured for toxicity versus the caf-
371 feine and the untreated N2a/A β PPsw cells, which
372 served as the control. The MTT assay showed no
373 significant difference from the control for toxicity as
374 compared to each concentration of THC and caffeine
375 administer suggesting THC and caffeine lack toxicity
376 to the cells at each concentration assayed (Fig. 3B).

377 *THC can inhibit A β_{40} aggregation as shown by*
378 *ThT assay and western blot*

379 The ThT assay was to exhibit the direct interaction
380 THC has with $A\beta$ demonstrates that as the concen-




Fig. 2. $A\beta_{40}$ (pg/ml) concentration in N2a/A β PPsw cells at various drug concentrations among groups. Treatment with both THC and caffeine resulted in a dose-dependent decrease in $A\beta_{40}$ concentration. However, no synergistic effect was observed.

381 tration of THC added to the assay was increased, the
382 intensity of fluorescence in $A\beta$ decreased. This data
383 suggests that $A\beta$ peptide directly binds to THC and
384 prevents the uptake of fluorescence (Fig. 4A). More-
385 over, our lab performed an additional ELISA assay to
386 confirm that the interaction of the $A\beta$ peptide with
387 THC did not shield amino acids 1–10, the major B-cell
388 epitope [40] (Fig. 4B). There is no significant difference
389 in absorbance at each concentration of THC, indicating
390 that at each concentration of THC the $A\beta$ antibodies
391 were able to bind with equal distribution and affinity.




Fig. 3. (A) $A\beta_{40}$ (pg/ml) concentration N2a/A β PPsw cells treated with THC, as well as the $A\beta_{40}$ (pg/ml) concentration of N2a/A β PPsw cells treated with THC twice, 24 hours apart. The number of treatments has shown to decrease the concentration of $A\beta_{40}$ (pg/ml), (B) This shows the data obtained from the reduction of MTT at different concentrations of THC versus the different concentration of caffeine. Untreated N2a/A β PPsw cells were also assayed to compare with the MTT reduction of N2a/A β PPsw cells treated with THC and caffeine at different concentrations.




Fig. 4. (A) ThT assay measuring the fluorescence of Thioflavin T which binds to β -sheet structure of $A\beta$ aggregation. With addition of THC, dose-dependent decreases in intensity of fluorescence indicates THC directly interferes with the binding of ThT to $A\beta$ peptide. (B) THC incubated with $A\beta$ peptide to determine the occurrence of THC interference with the major B cell epitope. No identified interference was observed at each increasing concentration of THC.




Fig. 5. (A) Polyacrylamide gel from a western blot indicating the concentration of aggregated $A\beta$ peptide with and without the treatment of THC at various concentrations. Groups: 1: Aggregation control; 2: THC 100 nM; 3: THC 10 nM; 4: THC 1 nM, (B) THC anti-aggregation assay gel quantification indicating the relative percent monomeric $A\beta$.




Fig. 6. ELISA assay elucidating a possible mechanism through which THC functions to decrease the synthesis of A β in N2a/A β PPSwe cells. A β level increases at 36 hours and reaches its peak level at 48 hours. Follow this mark; it then starts decreasing at 60 hours. The drug treatment benefit time is seen at 36 hours and last to 48 hours (the best window time). THC can significantly lower A β and this function can be partially blocked by CB1 antagonist Rimon at 10⁻⁴ M. However, inhibition function is lost at 10⁻⁷ M.

Therefore, we can postulate that THC's direct interaction with the A β peptide will not dampen an immune response to clear the A β peptide.

Further analysis with western blot was performed measuring the anti-aggregation properties of THC with A β peptide. At each increasing concentration of THC, a higher relative % of A β monomer was observed correlating with a lower intensity of aggregated A β peptide. This data suggests the direct interaction of THC with A β peptide and its ability to bind to the peptide and inhibit aggregation (Fig. 5A,B).

CB1 receptor antagonist can partially rescue A β level inhibited by THC

An ELISA was performed to determine the mechanism of THC in supporting the reduction of A β in N2a/A β PPSwe cells. A known inhibitor of the CB1 receptor, rimonabant, was mixed with THC at different concentrations. Untreated N2a/A β PPSwe cell A β concentrations were used as a control. It was noted that a dose dependent increase in A β was observed as the concentration of the inhibitor was increased. A time dependent effect of the inhibitor was also witnessed as the assay was repeated at the 12-, 36-, 48-, 60-, and 72-hour mark (Fig. 6). Due to increasing A β concentrations as the inhibitor concentration is increased, this suggests that THC partially functions through the CB1 receptor to mediate the synthesis of A β . The RT-PCR results for CB1 receptor expression level showed that

there is no significant upregulation by THC to CB1 receptor (data not shown).

THC can inhibit total GSK-3 β and phosphorylated GSK-3 β (pGSK-3 β) production

The western blot assay performed to examine the effect of THC on GSK-3 β exhibits a dose-dependent decrease in GSK-3 β . β -actin, a housekeeping gene, was used as a control to indicate that GSK-3 β was expressed at a constant rate and that the changes in intensity are not related to the change in expression amount. As shown in Fig. 7A-D, this data suggests that THC is efficacious in modulating and ameliorating the expression of GSK-3 β and could decrease neuronal apoptosis by down regulating GSK-3 β .

THC can inhibit phosphorylated (pTau) production, but not affect A β PP production

We detected pTau and A β PP levels among different treatment conditions. THC can lower pTau expression level with dose-dependent administration, but we did not see the differences in A β PP levels detected with 6E10 antibody (Fig. 8A-F).

THC can enhance mitochondrial function but will not interfere with melatonin's enhancement of the mitochondria

Isolated mitochondria from N2a/A β PPSwe cells showed higher oxygen utilization when treated with THC. When combined with melatonin, the function of the mitochondria is not altered (Fig. 9A, B).

DISCUSSION

Advances in therapeutics to prevent AD, or delay the progression, are currently being made. Recent research has shown caffeine and coffee are effective in limiting cognitive impairment and AD pathology in the transgenic mouse model by lowering brain A β levels, which are thought to be central to the pathogenesis of AD [41]. Similarly, the current study shows the *in vitro* anti-A β activity of caffeine, and of another naturally occurring compound, THC.

N2a/A β PPSwe cells were incubated separately with various concentrations of caffeine, melatonin, and THC. The relative anti-A β effect of THC was observed to increase in a time dependent manner. A dose-dependent decrease in A β concentration was noticed at lower concentrations of THC, as compared to caf-




Fig. 7. (A) A western blot performed to determine the effects of THC on GSK-3β in N2a/AβPPswE. β-actin was used as a control to indicate that the expression rate was constant. The left indicator is molecular weight. Lane 1, 2, and 3 are β-actin level and lane 4, 5, and 6 are GSK-3β expression. 1 and 4 are cell controls, 2 and 5 are cells treated with 2.5 nM THC, and lane 3 and 6 are cells treated with 0.25 nM THC. THC can inhibit GSK-3β level at 2.4 nM concentration, (B) Graph representing the expression decrease in GSK-3β in a dose-dependent manner by using β-actin to obtain a value for the ratio of expressed GSK-3β. As shown in the bar graph, the total GSK-3β decrease after using β-actin standardized protein loading, (C) GSK-3β expression in N2a/AβPPswE treated with different drugs: Cells were plated in 6 well plate for overnight and then drugs were added into each designated wells in duplicate. Cells were lysed after 36 hours incubation. Proteins were loaded onto SDS-page gel and then blotted with each antibody after transfer onto PVDF membrane. Groups are: CTRL, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (D) Expression of pGSK-3β following melatonin and THC treatment in N2a/AβPPswE cells. *The same batch protein samples were used in this test as in Fig. 7C. Bands were quantified. One-way ANOVA was applied to the data. *p* < 0.05 when compared with control group. ***p* < 0.01 when compared with control group. Groups are: Ctrl, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM.

444 feine. Further evidence shows that N2a/AβPPswE
 445 cells, treated twice with THC, show an even greater
 446 reduction in Aβ levels at slightly higher concentra-
 447 tions. Although it might have been predicted that
 448 caffeine and THC may function in a synergistic effect
 449 to reduce the Aβ load in N2a/AβPPswE cells, no syn-
 450 ergy was observed.

The MTT assay confirmed that cells treated at effi-
 cacious concentration of THC showed no toxicity,
 suggesting such a treatment to be safe and effective
 for further experimentation in the AD animal model.
 However, valid arguments have transpired in recent
 times regarding the concern for acute and long-term
 memory impairment with the use of THC. It has been

451
 452
 453
 454
 455
 456
 457




Fig. 8. (A) AβPP expression in N2a/AβPPswe treated with different drugs. The sample protein samples as in Fig. 7C were used for western blotting assay. 6E10 anti-Aβ antibody was used to detect AβPP and β-Actin was detected as protein loading control. Groups are: CTRL, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (B) Quantification result of AβPP in western blotting: We used quantification method to further compare the differences among drug treatment to AβPP level. There are no statistical significant differences among all treatment ($p > 0.05$). This data indicates that THC did not change AβPP expression level. Groups are: Ctrl, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (C) Tau expression in N2a/AβPPswe treated with different drugs. The sample protein samples as in Fig. 7C were used for western blotting assay. Anti-Tau and pTau antibodies were used to detect AβPP and β-Actin was detected as protein loading control. Groups are: CTRL, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (D) No significant difference of pTau expression shown among six groups in N2a/AβPPswe cells. THC treatment has no function to pTau expression. Groups are: Ctrl, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (E) Expression of pTau/Tau following melatonin and THC treatment in N2a/AβPPswe cells. ⁺ $p < 0.05$ when compared with THC 25 nM and THC 0.25 nM groups. ^{*} $p < 0.01$ when compared with THC 25 nM, THC 2.5 nM, and THC 0.25 nM groups. [#] $p < 0.05$ when compared with THC 25 nM and THC 2.5 nM groups. Groups are: Ctrl, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM, (F) Expression of tau following melatonin and THC treatment in N2a/AβPPswe cells. ⁺ $p < 0.05$ when compared with the THC 25 nM, THC 2.5 nM, and THC 0.25 nM groups. ^{**} $p < 0.01$ when compared with the THC 25 nM, THC 2.5 nM, and THC 0.25 nM groups. [#] $p < 0.05$ when compared with THC 2.5 nM group. Groups are: Ctrl, Control; M1T2, 10⁻⁵ M Melatonin + 2.5 nM THC; M2T2, 10⁻⁶ M Melatonin + 2.5 nM THC; T1, THC 25 nM; T2, THC 2.5 nM; T3, THC 0.25 nM.




Fig. 9. (A) The enhancement of mitochondria function to cells treated with different: N2a/AβPPswe cells were cultured in 10 cm tissue culture plate and then treated with drugs for 36 hours and mitochondria were harvested and tested for their ability of using oxygen utilization. Ctrl, Control; MIT2, 10^{-5} M Melatonin + 2.5 nM THC; M2T2, 10^{-6} M Melatonin + 2.5 nM THC; T2, THC 2.5 nM; M1, 10^{-5} M Melatonin. (B) The enhancement of mitochondria function to cells treated with different: N2a/AβPPswe cells were cultured in 10 cm tissue culture plate and then treated with drugs for 36 hours and mitochondria were harvested and tested for their ability of using oxygen utilization. Ctrl, Control; MIT2, 10^{-5} M Melatonin + 2.5 nM THC; M2T2, 10^{-6} M Melatonin + 2.5 nM THC; T2, THC 2.5 nM; M1, 10^{-5} M Melatonin.

Table 1

Difference and percent decrease of Aβ₄₀ (pg/ml) in THC treated cells at 2.5 μg/ml compared with the control at different time points

Time Point	6 h	24 h	48 h
Control	1064.025	5303	5935.525
THC 2.5 μg/ml	965.827	3648.975	2894.175
Percentage of decreased Aβ ₄₀	9.23%	31.19%	51.24%

shown that memory impairment was identified in rats treated with THC [42]. It should be clear, however, that the memory impairment observed occurred at concentrations more than a thousand times higher than what is presented here as a beneficial treatment in AD model N2a/AβPPswe cells. The concentrations used in the study are considered to be extremely low, as the concentrations that we focused on in the study were from 2.5 nM of THC down to 0.25 nM of THC. Although some studies with ultra-low doses of THC have indicated neurotoxic roles [42], newer research shows a neuroprotective role and actually promotes elevation of phosphorylated cAMP response element-binding protein (pCREB) by increasing the levels of brain-derived neurotrophic factor in the frontal cortex [43]. Furthermore, the dosing used in our study is a lower concentration than that in the aforementioned research. Therefore, we believe that THC has a therapeutic value, and that at low enough doses, the potential benefits strongly prevail over the risks associated with THC and memory impairment.

In addition to the Aβ concentration suppression, benefits of THC, analyzed with a western blot and ThT assay, confirmed anti-Aβ aggregate properties by a dose-dependent decrease in fluorescence uptake,

and a decrease in intensity of aggregated Aβ in a dose-dependent manner. The positive results suggest possible intermolecular force interactions, preventing the molecular aggregation of Aβ peptides. The conducted ELISA, to ensure the intermolecular interaction of THC with Aβ did not block the major B-cell epitope, showed no interference with antibody binding, which indicated that regardless of the molecular interaction of THC with Aβ, an immune response should not be inhibited.

One pathway in which THC function was shown through the cannabinoid receptor inhibition with rimonabant. The dose- and time-dependent increase of Aβ with respect to CB₁ inhibition was noted. It is likely that the time deference was observed due to the slow interaction of rimonabant with the CB₁ receptor. However, the difference in Aβ concentration becomes more evident at the later time points. Lastly, we showed a dose-dependent decrease in GSK-3β expression influenced by THC.

To date, no Aβ specific therapeutic options for AD have been approved. While progression is being made in this field, rigorous efforts focus on developing compounds that can address or possess the inhibition of Aβ synthesis and anti-Aβ aggregation properties or characteristics that down regulate GSK-3β and pGSK-3β. Our results demonstrate that THC possesses all of the above mentioned properties. All of these areas address major etiological characteristics of AD. GSK-3β, pGSK-3β, and Aβ-plaque brain concentrations, the hallmark of AD, are major targets for current AD research. Furthermore, we have shown that THC functions are pathway dependent of the endoge-

nous cannabinoid CB₁ receptor recently discovered to possibly function in AD disease modulation by suppressing microglial activation upon receptor interaction. Notwithstanding, it should also be noted that low doses of THC are used to address the above mentioned targets, thus avoiding risks induced by THC associated with memory impairment and risks associated with toxicity. In addition, we also discovered that low doses of THC can also enhance mitochondria function and has no negative drug interactions to melatonin, a potential therapeutic for AD.

Here we have presented a promising compound that addresses many major targets for AD therapeutics currently being research. We have shown THC, at an extremely low dose level (2.5 nM), has the proclivity to slow or halt AD progression by dampening the synthesis of the major pathological marker of AD, A β . Also, our lab has elucidated a potential mechanism responsible for the anti-pathological properties of THC with respect to AD. Furthermore, we have clearly exhibited lack of toxicity at low concentrations of both THC and caffeine individually. In conclusion, we believe the multifaceted functions of THC will ultimately decrease downstream tau hyperphosphorylation and neuronal death thereby halting or slowing the progression of this devastating disease.

DISCLOSURE STATEMENT

Authors' disclosures available online (<http://www.j-alz.com/disclosures/view.php?id=2309>).

REFERENCES

- [1] Alzheimer's, Association (2012) 2012 Alzheimer's disease facts and figures. *Alzheimers Dement* **8**, 131-168.
- [2] Saxena U (2011) Bioenergetics breakdown in Alzheimer's disease: Targets for new therapies. *Int J Physiol Pathophysiol Pharmacol* **3**, 133-139.
- [3] Götz J, Eckert A, Matamalas M, Ittner LM, Liu X (2011) Modes of A β toxicity in Alzheimer's disease. *Cell Mol Life Sci* **68**, 3359-3375.
- [4] Brookmeyer R, Johnson E, Ziegler-Graham K, Arrighi HM (2007) Forecasting the global burden of Alzheimer's disease. *Alzheimers Dement* **3**, 186-191.
- [5] Chiti F, Dobson CM (2006) Protein misfolding, functional amyloid, and human disease. *Annu Rev Biochem* **75**, 333-366.
- [6] Campbell V, Gowran A (2007) Alzheimer's disease; taking the edge off with cannabinoids? *Br J Pharmacol* **152**, 655-662.
- [7] Rich JB, Rasmussen DX, Folstein MF, Carson KA, Kawas C, Brandt J (1995) Nonsteroidal anti-inflammatory drugs in Alzheimer's disease. *Neurology* **45**, 51-55.
- [8] Octave JN (1995) The amyloid peptide and its precursor in Alzheimer's disease. *Rev Neurosci* **6**, 287-316.
- [9] Reitz C, Brayne C, Mayeux R (2011) Epidemiology of Alzheimer disease. *Nat Rev Neurol* **7**, 137-152.
- [10] Pillay N, Kellaway L, Kotwal G (2004) Molecular mechanisms, emerging etiological insights and models to test potential therapeutic interventions in Alzheimer's disease. *Curr Alzheimer Res* **1**, 295-306.
- [11] Hooper C, Killick R, Lovestone S (2008) The GSK3 hypothesis of Alzheimer's disease. *J Neurochem* **104**, 1433-1439.
- [12] Proctor CJ, Gray DA (2010) GSK3 and p53 - is there a link in Alzheimer's disease? *Mol Neurodegener* **5**, 7.
- [13] Lovestone S, Reynolds CH, Lätimer D, Davis DR, Ander-ton BH, Gallo JM, Hanger D, Mulot S, Marquardt B, Stabel S, et al. (1994) Alzheimer's disease-like phosphorylation of the microtubule-associated protein tau by glycogen synthase kinase-3 in transfected mammalian cells. *Curr Biol* **4**, 1077-1086.
- [14] Ishiguro K, Omori A, Takamatsu M, Sato K, Arioka M, Uchida T, Imahori K (1992) Phosphorylation sites on tau by tau protein kinase I, a bovine derived kinase generating an epitope of paired helical filaments. *Neurosci Lett* **148**, 202-206.
- [15] Hanger DP, Hughes K, Woodgett JR, Brion JP, Anderton BH (1992) Glycogen synthase kinase-3 induces Alzheimer's disease-like phosphorylation of tau: Generation of paired helical filament epitopes and neuronal localisation of the kinase. *Neurosci Lett* **147**, 58-62.
- [16] Cho JH, Johnson GV (2003) Glycogen synthase kinase 3beta phosphorylates tau at both primed and unprimed sites. Differential impact on microtubule binding. *J Biol Chem* **278**, 187-193.
- [17] Asuni AA, Hooper C, Reynolds CH, Lovestone S, Anderton BH, Killick R (2006) GSK3alpha exhibits beta-catenin and tau directed kinase activities that are modulated by Wnt In. *Eur J Neurosci* **24**, 3387-3392.
- [18] Hernandez F, Lucas JJ, Avila J (2013) GSK3 and tau: Two convergence points in Alzheimer's disease. *J Alzheimers Dis* **33**(Suppl 1), S141-S144.
- [19] Engel T, Goni-Oliver P, Lucas JJ, Avila J, Hernandez F (2006) Chronic lithium administration to FTDP-17 tau and GSK-3beta overexpressing mice prevents tau hyperphosphorylation and neurofibrillary tangle formation, but pre-formed neurofibrillary tangles do not revert. *J Neurochem* **99**, 1445-1455.
- [20] Phiel CJ, Wilson CA, Lee VM, Klein PS (2003) GSK-3alpha regulates production of Alzheimer's disease amyloid-beta peptides. *Nature* **423**, 435-439.
- [21] Sun X, Sato S, Murayama O, Murayama M, Park JM, Yamaguchi H, Takashima A (2002) Lithium inhibits amyloid secretion in COS7 cells transfected with amyloid precursor protein C100. *Neurosci Lett* **321**, 61-64.
- [22] DaRocha-Souto B, Coma M, Perez-Nievas BG, Scotton TC, Siao M, Sanchez-Ferrer P, Hashimoto T, Fan Z, Hudry E, Barroeta I, Sereno L, Rodriguez M, Sanchez MB, Hyman BT, Gomez-Isla T (2012) Activation of glycogen synthase kinase-3 beta mediates beta-amyloid induced neuritic damage in Alzheimer's disease. *Neurobiol Dis* **45**, 425-437.
- [23] Jackson SJ, Diemel LT, Pryce G, Baker D (2005) Cannabinoids and neuroprotection in CNS inflammatory disease. *J Neurol Sci* **233**, 21-25.
- [24] Riedel G, Davies SN (2005) Cannabinoid function in learning, memory and plasticity. *Handb Exp Pharmacol* **445**-477.
- [25] Van Sickle MD, Duncan M, Kingsley PJ, Mouihate A, Urbani P, Mackie K, Stella N, Makriyannis A, Piomelli D, Davison JS, Marnett LJ, Di Marzo V, Pittman QJ, Patel KD, Sharkey KA (2005) Identification and functional characterization of brainstem cannabinoid CB2 receptors. *Science* **310**, 329-332.
- [26] Ashton JC, Friberg D, Darlington CL, Smith PF (2006) Expression of the cannabinoid CB2 receptor in the rat

- 633 cerebellum: An immunohistochemical study. *Neurosci Lett*
634 **396**, 113-116.
- 635 [27] Nunez E, Benito C, Pazos MR, Barbachano A, Fajardo O,
636 Gonzalez S, Tolon RM, Romero J (2004) Cannabinoid CB2
637 receptors are expressed by perivascular microglial cells in the
638 human brain: An immunohistochemical study. *Synapse* **53**,
639 208-213.
- 640 [28] Marchalant Y, Brothers H, Norman G, Karelina K, DeVries
641 A, Wenk G (2009) Cannabinoids attenuate the effects of aging
642 upon neuroinflammation and neurogenesis. *Neurobiol Dis* **34**,
643 300-307.
- 644 [29] Ramírez B, Blázquez C, Gómez del Pulgar T, Guzmán M, de
645 Ceballos M (2005) Prevention of Alzheimer's disease pathol-
646 ogy by cannabinoids: Neuroprotection mediated by blockade
647 of microglial activation. *J Neurosci* **25**, 1904-1913.
- 648 [30] Benito C, Nunez E, Tolon RM, Carrier EJ, Rabano A, Hillard
649 CJ, Romero J (2003) Cannabinoid CB2 receptors and fatty
650 acid amide hydrolase are selectively overexpressed in neu-
651 ritic plaque-associated glia in Alzheimer's disease brains. *J*
652 *Neurosci* **23**, 11136-11141.
- 653 [31] Martín-Moreno AM, Reigada D, Ramírez BG, Mechoulam
654 R, Innamorato N, Cuadrado A, de Ceballos ML (2011)
655 Cannabidiol and other cannabinoids reduce microglial acti-
656 vation *in vitro* and *in vivo*: Relevance to Alzheimer's disease.
657 *Mol Pharmacol* **79**, 964-973.
- 658 [32] Bisogno T, Di Marzo V (2010) Cannabinoid receptors and
659 endocannabinoids: Role in neuroinflammatory and neurode-
660 generative disorders. *CNS Neurol Disord Drug Targets* **9**,
661 564-573.
- 662 [33] Piomelli D (2003) The molecular logic of endocannabinoid
663 signalling. *Nat Rev Neurosci* **4**, 873-884.
- 664 [34] Volicer L, Stelly M, Morris J, McLaughlin J, Volicer BJ (1997)
665 Effects of dronabinol on anorexia and disturbed behavior in
666 patients with Alzheimer's disease. *Int J Geriatr Psychiatry*
667 **12**, 913-919.
- 668 [35] Walther S, Mahlberg R, Eichmann U, Kunz D (2006) Delta-
669 9-tetrahydrocannabinol for nighttime agitation in severe
670 dementia. *Psychopharmacology (Berl)* **185**, 524-528.
- [36] Mishra S, Mishra M, Seth P, Sharma SK (2011) Tetrahy-
671 drocurcumin confers protection against amyloid β -induced
672 toxicity. *Neuroreport* **22**, 23-27. 673
- [37] Cao C, Loewenstein DA, Lin X, Zhang C, Wang L, Duara R,
674 Wu Y, Giannini A, Bai G, Cai J, Greig M, Schofield E, Ashok
675 R, Small B, Potter H, Arendash GW (2012) High blood caf-
676 feine levels in MCI linked to lack of progression to dementia.
677 *J Alzheimers Dis* **30**, 559-572. 678
- [38] Dragicevic N, Copes N, O'Neal-Moffitt G, Jin J, Buzzeo R,
679 Mamcarz M, Tan J, Cao C, Olcese JM, Arendash GW, Brad-
680 shaw PC (2011) Melatonin treatment restores mitochondrial
681 function in Alzheimer's mice: A mitochondrial protective role
682 of melatonin membrane receptor signaling. *J Pineal Res* **51**,
683 75-86. 684
- [39] Cao C, Cirrito J, Lin X, Wang L, Verges D, Dickson A, Mam-
685 carz M, Zhang C, Mori T, Arendash G, Holtzman D, Potter H
686 (2009) Caffeine suppresses amyloid-beta levels in plasma and
687 brain of Alzheimer's disease transgenic mice. *J Alzheimers*
688 *Dis* **17**, 681-697. 689
- [40] Agadjanyan MG, Ghochikyan A, Petrushina I, Vasilevko V,
690 Movsesyan N, Mkrtychyan M, Saing T, Cribbs DH (2005)
691 Prototype Alzheimer's disease vaccine using the immun-
692 odominant B cell epitope from beta-amyloid and promiscuous
693 T cell epitope pan HLA DR-binding peptide. *J Immunol* **174**,
694 1580-1586. 695
- [41] Arendash G, Cao C (2010) Caffeine and coffee as therapeutics
696 against Alzheimer's disease. *J Alzheimers Dis* **20**(Suppl 1),
697 S117-S126. 698
- [42] Nakamura EM, da Silva EA, Concilio GV, Wilkinson DA,
699 Masur J (1991) Reversible effects of acute and long-term
700 administration of delta-9-tetrahydrocannabinol (THC) on
701 memory in the rat. *Drug Alcohol Depend* **28**, 167-175. 702